

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/334989163>

ALOE VERA: AN ASSURED WEIGHT LOSS DIET –AN APPROACH TOWARD IMPROVING THE JUICE PALATABILITY AND IN SILICO ANALYSIS

Article in *Asian Journal of Pharmaceutical and Clinical Research* · May 2019

DOI: 10.22159/ajpcr.2019.v12i6.32380

CITATIONS

0

READS

253

3 authors:



Disalva Xavier
Vels University

5 PUBLICATIONS 67 CITATIONS

[SEE PROFILE](#)



Sharanya Manoharan
Vels University

29 PUBLICATIONS 42 CITATIONS

[SEE PROFILE](#)



S.Radha Mahendran
Vels University

89 PUBLICATIONS 108 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



English Language Teaching Methodology [View project](#)



INSILICO STUDIES ON VARIOUS NATURAL INHIBITORS [View project](#)

ALOE VERA: AN ASSURED WEIGHT LOSS DIET – AN APPROACH TOWARD IMPROVING THE JUICE PALATABILITY AND *IN SILICO* ANALYSIS

DISALVA X¹, SHARANYA M^{2*}, MAHENDRAN RADHA²

¹Department of Hotel and Catering Management, Vels Institute of Science, Technology and Advanced Studies, Chennai, Tamil Nadu, India.

²Department of Bioinformatics, Vels Institute of Science, Technology and Advanced Studies, Chennai, Tamil Nadu, India.

Email: sharanya.bioinfo@gmail.com

Received: 03 February 2019, Revised and Accepted: 16 April 2019

ABSTRACT

Objective: Obesity is an epidemic disease act as causative for global death. The principal aim of this study is to create an obesity treatment beverage that is palatable and readily acceptable to the public. *Aloe vera* juice is one such drink known to assist in reducing obesity. This juice is sold at several places in Chennai and more easily available for morning walkers along the Marina beach. Although known for its benefits in promoting a slim and fit physique, its taste is less palatable. The objective of this study is to develop a concoction that will make the *A. vera* juice tasty without compromising its nutritive value.

Materials and Methods: The taste is achieved by the addition of various fruit juices such as orange, lime, sweet lime, muskmelon, and pineapple with honey and stevia for sweetening. The weight loss ability of the prepared drinks is evaluated using pancreatic lipase inhibitory and the presence of phytochemicals. To the credit, the present study also determines the efficiency of *A. vera* compounds for its antiobesity property through *in silico* techniques. The significant interaction exhibited by the compounds with the antiobesity target inositol hexakisphosphate kinase 1 (IP6K1) was discussed.

Results: The recipe B containing orange juice: *A. vera* juice: stevia in the ratio 3:3:1 had good taste and the significant lipase activity. The phytochemicals present in the *A. vera* are tannins, flavonoids, alkaloids, steroids, and polyphenols, and these phytochemicals are observed having significant interaction with protein IP6K1. Aloin A and aloe emodin had significant Glide score and interactions with active site residues.

Conclusion: Natural herbal products for weight reduction may be effective in the treatment of obesity and associated disorders. The potential lipase inhibition activity of juice may be due to the presence of various phytochemicals such as flavonoids, polyphenol etc. in the *Aloe vera*.

Keywords: Obesity, *Aloe vera*, Weight loss, *Aloe vera* juice, *In silico* analysis, Docking studies, Inositol hexakisphosphate kinase 1.

© 2019 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>) DOI: <http://dx.doi.org/10.22159/ajpcr.2019.v12i6.32380>

INTRODUCTION

In the present era, obesity is one among the causes for global death and other diseases. It is an epidemic disease with increased comorbidity for Type 2 diabetes, coronary heart disease, stroke, fatty liver, neurodegeneration, cerebrovascular disease, congestive heart failure, hypertension, lipoprotein abnormalities, and several diseases [1]. Obesity is an abnormal or excessive fat accumulation to the body and could be explained as a body mass index, kg/m². More recently, Guthold *et al.* [2] published that 28% or 1.4 billion people are physically inactive which increases the risk for obesity and the development of non-communicable diseases. Despite physical inactivity, excess intake of food, especially processed junk foods, endocrine disorders, genes, medications, and mental disorders also have its role in the cause of obesity [3,4].

Aloe vera, a common medicinal, readily available, cactus-like plant has several medicinal applications in treating many ailments. Christaki and Florou-Paneri [5] reviewed the worth of *A. vera* under the categories cosmetic, medicinal, food, and nutrition. This plant being used since Rigvedic times in India, being reported for its medicinal property such as treating wounds and burns, protection for skin from X-rays, lung cancer, intestinal problems, reduces the blood sugar level in diabetic patients and low-density lipoprotein, increase the high-density lipoprotein, improves the immune system as well as to fight the acquired immune deficiency syndrome (AIDS), and allergies [6,7]. *A. vera* gained its industrial importance and the value for their products roughly calculated around \$125 million and the recent thrust in utilizing herbal medicines have enlightened its significance in the industrial

development [6]. A report was found that Americans spend 40 billion dollars on Aloe products as functional foods, drinks, and nutritional supplements [6]. Aloe juices are widely used as a health drink, soft drink, laxative drink, sherbet and along with several components such as lemon juice, electrolyte, soluble fiber, Vitamin B, amino acids and acetaminophen, other vegetables, yogurts, and cucumber [8]. Due to its multi-application, the Chinese called the plant as Elixir of youth. The intake of Aloe preparations as a whole leaf extract and as an inner fillet gel enhanced the uptake of water/fat-soluble vitamins in the body and it is the only supplement known for this ability [9]. The applications of Aloe in health are enormous and scientific records on its antiacid, antipeptic, gastroprotective, and antiulcer [10].

The emergence of bioinformatics has led the drug discovery and development process much easier, also cost effective and the existing high throughput sequencing techniques added its value. In addition, the juice prepared as a daily dietary ingredient, the present study focuses to study the potentials of *A. vera* compounds in interacting the antiobesity target inositol hexakisphosphate kinase 1 (IP6K1). The target chosen has significant functionality in regulating cell metabolism and survival [11]. The recent studies have found that deletion of adipocyte-specific IP6K1 modulates AMP-activated protein kinase-mediated adipocyte energy metabolism thereby regulates the fat accumulation in the body [12]. Secondary metabolites of plants have gained momentum in the process of drug discovery. Recently, arteether, galantamine, nitisinone, and tiotropium have been introduced in the US market [13]. Even more, the plant natural compounds were scientifically evaluated its efficiency against several dreadful diseases such as cancer, HIV/AIDS, Alzheimer's, and malaria [14].

MATERIALS AND METHODS

Beverage preparation

The beverage preparation consists of three steps which include juice processing of selected fruits for improving taste, *A. vera* juice processing and beverage preparation. The three citrus fruits juices of mosambi, orange, and lemon as well as the pineapple and muskmelon were chosen and added with natural sweeteners in the form of powder (stevia and honey). The *A. vera* leaves were bought from Pallavaram Market, Chennai. The juices of each fruit were prepared in the following methods. The skins of mosambi was removed and roughly blended to extract the juice and stored immediately since citrus fruits have a tendency to turn to bitter taste. In the case of orange and lemon, squeezing techniques were used to obtain the extract and mixed with 3/4th of water. In pineapple and muskmelon, the skin was removed, and the pulp was diced into pieces to prepare as juice. The prepared juices were kept aside and/or stored in the refrigerator until the *A. vera* leaves were extracted. *A. vera* leaves were peeled, diced, and washed several times to remove the stickiness and pulp was extracted by the blending method and processed into juice. Furthermore, it was filtered through a strainer and stored in clean glass jar. As a final step, the beverage was prepared by mixing together the 30 ml of each fruit juices with 30 ml of *A. vera* juice separately. The mixture was blended nicely using cocktail shaker with the addition of sweeteners. The recipe formulation was tabulated (Table 1).

Determination of phytochemicals and nutraceuticals

Phytochemical analysis

The phytochemical analysis was carried out for the food samples prepared for the presence of tannins, flavonoids, steroids, alkaloids, and polyphenols [15-17]. The samples were extracted with hydroalcoholic solvents (70%) (70 ml ethanol and 30 ml of water) for 24 h, and the filtrates were analyzed for phytochemicals.

Test for tannins

About 1 ml of the sample was boiled in 20 ml of water in a test tube and then filtered. A few drops of 0.1% ferric chloride was added and observed for brownish green or a blue-black coloration.

Test for flavonoids

About 5 ml of dilute ammonia solution were added to a portion of the aqueous filtrate of each plant extract followed by addition of concentrated H₂SO₄. A yellow coloration observed in each extract indicated the presence of flavonoids. The yellow coloration disappeared on standing.

Table 1: Composition of concoction

S. No.	Recipe formulation	Quantity of fruits juice (ml)	Quantity of <i>A. vera</i> juice (ml)	Quantity of stevia (g)
1	A	Lemon - 30	30	10
2	B	Orange - 30	30	10
3	C	Sweet Lime - 30	30	10
4	D	Pineapple - 30	30	10
5	E	Muskmelon - 30	30	10

A. vera: *Aloe vera*

Test for steroids

About 2 ml of acetic anhydride was added to 1 ml of an extract of each sample with 2 ml H₂SO₄. The color changed from violet to blue or green in some samples indicate the presence of steroids.

Test for alkaloids

Mayer's test - to a few (1) ml of the extract, a drop of Mayer's reagent was added by the side of the test tube. A creamy or white precipitate indicates positive.

Test for polyphenols

Ethanol of 4 ml was added to each extract (1 ml) and the resulting solution was transferred in test tubes and kept warm in a water bath for 15 min. Three drops of freshly prepared ferric cyanide solution were added to the extract solution. Formation of a blue-green color indicated the presence of polyphenols.

Determination of weight loss

The efficiency of weight loss was tested using pancreatic lipase inhibitory activity. The lipase enzyme was extracted from the pancreas of chicken (*Gallus domesticus*).

Extraction of lipase from chicken (*G. domesticus*) pancreas

The method was followed according to Choi *et al.* [18]. The pancreas of freshly slaughtered chicken was collected, washed thoroughly and placed in ice-cold sucrose solution (0.01 M). The pancreas was homogenized in 0.01 M sucrose and centrifuged. The supernatant solution was separated and subjected to ammonium sulfate precipitation (50% saturation). The obtained white pellets after centrifugation were dissolved in sucrose solution and again saturated with 50% ammonium sulfate and centrifuged. Finally, pellets were used as enzyme source by dissolving in phosphate buffer.

Pancreatic lipase inhibitory activity

The lipase inhibitory activity was determined using the titration method [19]. Different concentrations of juice samples were taken in the concentration of 50, 100, and 150 ml and further mixed with 8 ml of olive oil, 0.4 ml phosphate buffer, and 1 ml of chicken pancreatic lipase. The mixture was incubated for 60 min. The reaction was terminated by the addition of 1.5 ml of a solution containing acetone and 95% ethanol (1:1). The appearance of pink color from yellow color indicates the liberated fatty acids, which was determined by titrating the solution against 0.02 M sodium hydroxide (standardized by 0.01 M oxalic acid) using phenolphthalein as an indicator and the percentage inhibition of lipase activity was calculated using the following formula:

$$\text{Lipase inhibition percentage} = \frac{\text{Lipase activity before treatment} - \text{Lipase activity after treatment}}{\text{Lipase activity before treatment}} \times 100$$

In silico analysis

The protein sequence of antiobesity target IP6K1 was retrieved from UniProt database (Accession ID: Q92551) from the *Homo sapiens*,

Table 2: Phytochemicals analysis of different juices

S. No.	Juice varieties	Phytochemicals				
		Tannin	Flavonoids	Steroids	Alkaloids	Polyphenols
1	Lemon with <i>A. vera</i>	+	++	+	+	++
2	Orange with <i>A. vera</i>	+	++	+	+	++
3	Sweet lime with <i>A. vera</i>	+	++	++	+	++
4	Pine apple with <i>A. vera</i>	+	++	+	+	++
5	Muskmelon with <i>A. vera</i>	+	++	+	+	++

Note: + indicates presence, ++ indicates high concentration, *A. vera*: *Aloe vera*

which contains 441 amino acids and the mass of 50,236 Da. Further, the retrieved sequence was subjected for alignment using the protein-protein basic local alignment search tool (BLAST) against the protein structure database (Protein Data Bank [PDB]), to identify the structure of similar proteins for protein modeling. The structure prediction was carried out using the online Tool I-Tasser, and was an automated structure prediction tools from the structure-based function annotation (<https://zhanglab.ccmb.med.umich.edu/I-TASSER/>). The quality of the modeled structure was studied from the Ramachandran Plot obtained using the tool, RamPage (<http://mordred.bioc.cam.ac.uk/~rapper/rampage.php>). Before the energy of the structure was minimized using Swiss PDB viewer. The active site pocket for the modeled protein structure was predicted using SCFbio (<http://www.scfbio-iitd.res.in/dock/ActiveSite.jsp>). Plant compounds emodin, anthracene, aloemodin, chrysophanic acid, anthranol, aloin B, aloetic acid, aloin, aloin A, and the Food and Drug Administration approved drugs for obesity, orlistat, and phentermine in three dimensional. sdf file format was retrieved from the PubChem database (<https://pubchem.ncbi.nlm.nih.gov/>). Further, the docking studies were carried out using Glide module of Schrodinger software to analyze the docking score, and the interactions with specific residues along with the length of bond formation were visualized in PyMol software.

RESULTS AND DISCUSSION

Functional foods are defined as foods that contain bio-active ingredients thought to enhance health and fitness. The active ingredients are phytochemicals, such as lycopene in tomatoes, allicin in garlic, or isoflavones in soybeans. These phytochemicals, also called "nutraceuticals," may be extracted and consumed as supplements, or may have therapeutic value when consumed in whole food. Majority of foods, such as whole grains, beans, fruits, vegetables, and herbs contain phytochemicals of nutraceutical importance. These phytochemicals, either alone and/or in combination, have tremendous therapeutic potential in curing various ailments including cancer, diabetic, obesity, arthritis, ulcer, cardiovascular diseases, and hypertension [20]. Natural products provide a vast pool of pancreatic lipase inhibitors [21]. A wide variety of plant products such as saponins, polyphenols, flavonoids, and caffeine possess lipase inhibitors. Natural products can play a safe and effective role with obesity especially those containing fibers, polyphenols, sterols, and alkaloids [22]. In the present study, tannin, flavonoids, steroids, alkaloid, and polyphenol were present in the tested sample (Table 2). The presence of these compounds indicates the positive effect on health.

Determination of weight loss

In the present era, human population faces obesity as the major problem. However, the number of agents and their products is utilized for weight loss; perhaps the side effects are the issues to be concerned. Hence, consistent and safe herbal product for weight reduction is the urgent need in developing countries. The developed juice varieties were observed for lipase inhibition and the results are discussed (Table 3).

Lipase inhibitors may be effectiveness in reducing dietary fat intake by reducing both the consumption and absorption of fat [23]. Dietary fat is absorbed by the intestine when it has been subjected to the action of pancreatic lipases. Pancreatic lipase is a key enzyme in dietary triacylglycerol absorption, hydrolyzing triacylglycerols to monoacylglycerols, and fatty acids. Digestion and absorption of dietary lipids by pancreatic lipase, a major source of excess calorie intake, can be targeted for the development of weight loss agents. Pancreatic lipase inhibition is one of the most widely studied mechanisms used to determine the potential efficacy of natural products as antiobesity agents. Hence, in the present study, lipase was isolated from the chicken pancreas and determined the inhibitory activity of pancreatic lipase when incubated with different concentrations from 50, 100, 150, and 150 ml of Lemon with *A. vera*, orange with *A. vera*, sweet lime on with *A. vera*, pineapple with *A. vera*, and muskmelon with *A. vera* juice. Many

fruits and herbal teas have been extensively studied for the pancreatic lipase inhibition due to the presence of polyphenols [24,25].

With the increase in the concentration of extracts, the higher inhibition of the enzyme was observed. The order of lipase inhibition activity was Lemon with *A. vera*>Pineapple with *A. vera*>Muskmelon with *A. vera*>Orange with *A. vera*>Sweet lime on with *A. vera*. Comparatively lemon with *A. vera* juice showed maximum inhibition against enzyme lipase whereas the lowest inhibition was sweet lime on with *A. vera* juice. Thus, an inhibitor of digestive lipase that helps to limit intestinal fat absorption could be proved as useful medication for the treatment of hyperlipidemia and holds great promise as a weight loss agent. Among the various juice, lemon with *A. vera* juice possesses potential lipase inhibitors than other juice.

In silico analysis

Protein structure modeling

The protein IP6K1 from *H. sapiens* contains 441 amino acids, and the sequence was subjected for BLAST analysis. The hits obtained have only 32% of similarity against the PDB database; however, for obtaining a qualified structure, the template should be selected on the basis of selection rule with the highest sequence similarity [26]. Since in the present analysis, the percentage obtained was 32%, the multiple template-based structure prediction was performed; multiple template selection would increase the model accuracy [27]. I-Tasser generated five models, each with a C-score of -2.48, -2.72, -2.90, -3.95, and -2.74, respectively, whereas the estimated TM-Score and RMSD were 0.43 ± 0.14 and $13.0 \pm 4.2 \text{ \AA}$, respectively. The modeled structure was energy minimized using Swiss PDB viewer of -14389.468 KJ/Mol and the structure is shown in Fig. 1. The active sites were determined as THR 101, ASP 106, THR 107, THR 108, GLU 109, ARG 110, GLU 111, GLN 112, PRO 113, ARG 114, ARG 115, LYS 116, SER 118, ARG 119, ARG 124, SER 125, GLY 126, SER 127, GLY 18, SER 129, ASP 130, GLU133, GLU134, SER 137, LEU138,

Table 3: Lipase activity of different juices

Name of the juice	Concentration (ml)			
	0.50	1.00	1.50	2.00
Lemon with <i>A. vera</i>	8.50	7.80	7.20	6.32
Orange with <i>A. vera</i>	9.50	8.60	8.10	7.80
Sweet lime with <i>A. vera</i>	10.20	9.70	8.50	8.10
Pineapple with <i>A. vera</i>	9.20	8.50	7.80	7.10
Muskmelon with <i>A. vera</i>	9.80	8.70	8.20	7.40

A. vera: Aloe vera




Fig. 1: Structure of modeled protein-inositol hexakisphosphate kinase 1

Table 4: Absorption, distribution, metabolism, and excretion properties of *A. vera* compounds

Compound name	Molecular weight	Donor HB	Acceptor HB	QPlogPo/w Predicted octanol/water partition coefficient	SASA	QPlog BB Predicted blood/blood partition coefficient	Human Oral Absorption
	130.0–725.0	0.0–6.0	2.0–20.0	–2.0–6.5	300.0–1000.0	–3.0–1.2	1, 2 or 3 for low, medium or high
Orlistat (3034010)	495.741	0.25	7.75	21.148	961.847	–2.678	1
Emodin (3220)	270.241	1	4.25	12.607	479.225	–1.541	3
Phentermine (4771)	149.235	2	1	8.782	383.538	0.487	3
Anthracene (8418)	178.233	0	0	7.45	404.551	0.192	1
Aloe-emodin (10207)	270.241	1	5.2	12.997	478.805	–1.594	3
Chrysophanic acid (10208)	254.242	0	3.5	10.583	466.991	–0.987	3
Anthranol (10731)	194.232	1	0.75	9.357	411.593	0.193	3
Aloin B (14989)	418.399	5	11.7	26.431	626.291	–2.823	1
Aloetic acid (5464178)	450.231	1	9.2	18.705	597.175	–4.481	1
Aloin A (12305761)	418.399	5	11.7	26.634	641.8	–3.064	1

HB: Hydrogen bond, QPlogPo/w: Octanol/water coefficient, SASA: Solvent accessibility of surface area, QPlogBB: Blood-brain barrier, *A. vera*: *Aloe vera*

Table 5: Docking studies for IP6K1 with *A. vera* compounds

Compound Name	G. score (Kcal/mol)	Interacting residues	Bond length (Å)	No. of hydrogen Bonds
Aloin A (12305761)	–7.46	Lys218 (H-O)	2.2	5
		Lys218 (H-O)	1.7	
		Arg115 (H-O)	2.7	
		Gln112 (H-O)	2.4	
		Gln112 (O-H)	2.2	
Aloin B (14989)	–7.41	His272 (N-H)	2.0	2
		Arg115 (H-O)	2.1	
Aloe-Emodin (10207)	–7.35	Arg22 (O-H)	2.1	7
		Lys189 (O-H)	1.8	
		Arg124 (H-O)	2.1	
		Arg124 (H-O)	2.1	
		Arg119 (H-O)	1.9	
		Arg119 (H-O)	2.0	
		Asp43 (O-H)	1.6	
Aloetic acid (5464178)	–6.63	Arg41 (H-O)	2.6	8
		Arg41 (H-O)	2.6	
		Arg119 (H-O)	2.3	
		Arg119 (H-O)	1.9	
		Arg124 (H-O)	1.9	
		Asp130 (O-H)	2.0	
		Arg115 (H-O)	2.1	
		Lys189 (O-H)	1.8	
Emodin (3220)	–6.33	Lys189 (O-H)	1.8	6
		Lys132 (H-O)	2.6	
		Asp130 (O-H)	1.9	
		Arg124 (H-O)	2.1	
		Arg124 (H-O)	2.2	
		Arg22 (O-H)	2.0	
		Lys218 (O-H)	2.1	
Chrysophanic acid (10208)	–6.22	Lys218 (O-H)	2.1	5
		Lys218 (O-H)	2.9	
		Gln112 (H-O)	2.7	
		Arg114 (H-O)	2.7	
		Phe217 (O-HH)	1.8	
Anthranol (10731)	–4.91	Lys218 (O-H)	1.9	1
Phentermine (4771)	–4.33	Asp43 (O-H)	1.6	1
Anthracene (8418)	–3.95	NIL	NIL	NIL
Orlistat (3034010)	–2.89	Tyr266 (H-O)	2.7	2
		Lys218 (O-H)	2.2	

IP6K1: Inositol hexakisphosphate kinase 1

VAL214, PHE217, LYS218, TYR219, PRO220, GLN264, TYR266, GLN267, ASP269, GLY271, TYR273, GLY341, LYS342, GLU343, ARG345, GLU347, and SER348.

Absorption, distribution, metabolism, and excretion (ADME) properties


The absorption, distribution, metabolism, and excretion properties were determined for the *A. vera* compounds and drug molecule orlistat (Table 4). All the compounds analyzed had higher log p value; however, violation of single Lipinski rule of five is accepted. The molecules phentermine, anthracene, anthranol had a less molecular weight <200, emodin, aloe-emodin, and chrysophanic acid had in the range of 200–300, whereas orlistat, aloin B, aloetic acid, and aloin A had >400. Donor and acceptor hydrogen bond are most important since those are involved in the formation of significant interaction between the compound and the protein. The compound anthracene had no ability to donate as well as to accept the hydrogen bonds. The surface area solvent accessible is important criteria which reveal the polar interaction and the solubility of the compounds in the water environment. The compounds studied in the present research indicated significant solvent accessibility of surface area value. The table also shows the blood/blood partition coefficient for the compounds which ranges between the –3.0 and 1.2 and the human oral absorption was determined to predict the ability of the compounds for oral intake. The drug orlistat had a low level of oral absorption whereas the compounds emodin, phentermine, aloe-emodin, chrysophanic acid, and anthranol had significantly high for oral absorption.

Docking analysis


The docking studies for the compounds and drug orlistat with the protein IP6K1 1 were determined using the software Glide, a Schrodinger suite. The Glide score (G. Score), interacting residues, bond length, and number of hydrogen bonds were tabulated (Table 5). The plant compounds had significant G. score value that the drug orlistat where the G. score was observed to be –2.89 Kcal/mol. Among the plant compounds, aloin A had significant G. score of –7.46 Kcal/mol followed by aloin B and aloe emodin of –7.41 and –7.35 Kcal/mol, respectively. The respective G. score of aloetic acid, emodin, and chrysophanic acid is –6.63, –6.33, and –6.22 Kcal/mol. The compounds anthranol and phentermine had G. score in the range of –4 and anthracene had –3.95 Kcal/mol. The interactions were higher to lower in the order of aloetic acid>aloe emodin>emodin>aloin A, chrysophanic acid, orlistat>aloin B>anthranol, phentermine.

The interactions with active sites include GLN 112, ARG 115, ARG 119, ARG 124, ASP 130, PHE 217, LYS 218, and TYR 266. Perhaps, apart


Table 6: Snapshots of docking complex of compounds with protein IP6K1



Emodin versus IP6K1




Phentermine versus IP6K1




Aloe-Emodin versus IP6K1




Chrysophanic acid versus IP6K1




Anthranol versus IP6K1




Aloin B versus IP6K1



Aloetic acid versus IP6K1



Aloin A versus IP6K1



Orlistat versus IP6K1

IP6K1: Inositol hexakisphosphate kinase 1

from active sites, the following are the residues had interactions with the plant compounds and orlistat, ARG22, ARG 41, ASP 43, ARG 114, LYS 132, and LYS 189. The interactions of each compound with the protein IP6K1 are shown in Table 6. The compound anthracene had no interactions as it has no hydrogen donor and hydrogen acceptor property as predicted in the ADME analysis.

The *in silico* studies indicated the efficiency of *A. vera* compounds in interacting with the target protein IP6K1 and the ADME properties also indicated its human oral absorption property. The plant is reported in cleansing the digestive system and encourages in relaxing constipation as well as increases the energy levels to maintain a healthy body weight [28].

CONCLUSION

Obesity is characterized as abnormal or excessive fat deposition in adipose tissue and other internal organs such as liver, heart, and skeletal muscle. It is a chronic disorder of carbohydrate and fat metabolism and poses a risk to the health and well-being of humans. Consistent and safe herbal product for weight reduction is a need in developing countries. Natural herbal products for weight reduction may be effective in the treatment of obesity and associated disorders. The potential lipase inhibition activity of juice may be due to the presence of various phytochemicals such as flavonoids, polyphenol etc. in the *Aloe vera*.

AUTHOR'S CONTRIBUTIONS

The author declares that this work was done by the author named in this article.

CONFLICTS OF INTEREST

No conflicts of interest are associated with this work.

REFERENCES

- Berthoud HR, Klein S. Advances in obesity: Causes, consequences, and therapy. *Gastroenterology* 2017;152:1635-7.
- Guthold R, Stevens GA, Riley LM, Bull FC. Worldwide trends in insufficient physical activity from 2001 to 2016: A pooled analysis of 358 population-based surveys with 1.9 million participants. *Lancet Glob Health* 2018;6:e1077-86.
- Karam JG, McFarlane SI. Secondary causes of obesity. *Therapy* 2007;4:641-50.
- Bollapragada MK, Shantaram M, Kumar RS. Obesity: Development, epidemiology, factors affecting, quantity, health hazards, management and natural treatment a review. *Int J Pharm Pharm Sci* 2017;9:12-26.
- Christaki EV, Florou-Paneri PC. *Aloe vera*: A plant for many uses. *J Food Agric Environ* 2010;8:245-9.
- Ahlawat KS, Khatkar BS. Processing, food applications and safety of *Aloe vera* products: A review. *J Food Sci Technol* 2011;48:525-33.
- Subhshis P, Dutta S, Chaudhuri TK, Bhattacharjee SB. Anti-inflammatory and protective properties of *Aloe vera* leaf leaf crude gel in carrageenan induced acute inflammatory rat models. *Int J Pharm Pharm Sci* 2014;6:368-71.
- Singh A, Singh AK. Optimization of processing variables for the preparation of herbal bread using *Aloe vera* gel. *J Food Sci Technol* 2009;46:335-8.
- Vinson JA, Al Kharrat H, Andreoli L. Effect of *Aloe vera* preparations on the human bioavailability of Vitamins C and E. *Phytomedicine* 2005;12:760-5.
- Borrelli F, Izzo AA. The plant kingdom as a source of anti-ulcer remedies. *Phytother Res* 2000;14:581-91.
- Chakraborty A, Koldobskiy MA, Bello NT, Maxwell M, Potter JJ, Juluri KR, et al. Inositol pyrophosphates inhibit akt signaling, thereby regulating insulin sensitivity and weight gain. *Cell* 2010;143:897-910.
- Zhu Q, Ghoshal S, Rodrigues A, Gao S, Asterian A, Kamenecka TM, et al. Adipocyte-specific deletion of ip6k1 reduces diet-induced obesity by enhancing AMPK-mediated thermogenesis. *J Clin Invest* 2016;126:4273-88.
- Balunas MJ, Kinghorn AD. Drug discovery from medicinal plants. *Life Sci* 2005;78:431-41.
- Liu J, Pei M, Zheng C, Li Y, Wang Y, Lu A, et al. A systems-pharmacology analysis of herbal medicines used in health improvement treatment: Predicting potential new drugs and targets. *Evid Based Complement Alternat Med* 2013;2013:938764.
- Sofowora A. *Medicinal Plants and Traditional Medicinal in Africa*. 2nd ed. Sunshine House, Ibadan, Nigeria: Spectrum Books Ltd.; 1993.
- Trease GE, Evans WC. *Pharmacognosy*. 13th ed. London: ELBS/Bailliere Tindall; 1989.
- Harborne JB. *Phytochemical methods*. London Chapman and Hall Ltd., 1984.
- Choi SJ, Hwang JM, Kim SI. A colorimetric microplate assay method for high throughput analysis of lipase activity. *J Biochem Mol Biol* 2003;36:417-20.
- Kekuda TR, Raghavendra HL, Mallikarjun N, Venugopal TM, Kumar HS. Elemental composition, anticarcinogenic, pancreatic lipase inhibitory and cytotoxic activity of *Artocarpus lakoocha* Roxb pericarp. *Int J Drug Dev Res* 2012;4:330-6.
- Prakash D, Gupta C, Sharma G. Importance of phytochemicals in nutraceuticals. *J Chin Med Res Dev* 2012;1:70-8.
- Birari RB, Bhutani KK. Pancreatic lipase inhibitors from natural sources: Unexplored potential. *Drug Discov Today* 2007;12:879-89.
- Shimoda H, Seki E, Aitani M. Inhibitory effect of green coffee bean extract on fat accumulation and body weight gain in mice. *BMC Complement Altern Med* 2006;6:9-13.
- Ellrichmann M, Kapelle M, Ritter PR, Holst JJ, Herzig KH, Schmidt WE, et al. Orlistat inhibition of intestinal lipase acutely increases appetite and attenuates postprandial glucagon-like peptide-1-(7-36)-amide-1, cholecystokinin, and peptide YY concentrations. *J Clin Endocrinol Metab* 2008;93:3995-8.
- Nakai M, Fukui Y, Asami S, Toyoda-Ono Y, Iwashita T, Shibata H, et al. Inhibitory effects of oolong tea polyphenols on pancreatic lipase *in vitro*. *J Agric Food Chem* 2005;53:4593-8.
- Li F, Li W, Fu H, Zhang Q, Koike K. Pancreatic lipase-inhibiting triterpenoid saponins from fruits of *Acanthopanax senticosus*. *Chem Pharm Bull (Tokyo)* 2007;55:1087-9.
- Fiser A. Template-based protein structure modeling. *Methods Mol Biol* 2010;673:73-94.
- Fernandez-Fuentes N, Madrid-Aliste CJ, Rai BK, Fajardo JE, Fiser A. M4T: A comparative protein structure modeling server. *Nucleic Acids Res* 2007;35:W363-8.
- Rajeswari R, Umadevi M, Rahale S, Pushpa R, Selvavenkadesh S, Kumar KP, et al. *Aloe vera*: The miracle its medicinal and traditional uses in India. *J Pharmacogn Phytochem* 2012;1:118-24.